

Anomalous metastability in a temperature-driven transition

Miguel Ibáñez Berganza¹, Pietro Coletti², Alberto Petri³

¹*IPCF-CNR, UOS Roma Kerberos and Dipartimento di Fisica,*

Università “La Sapienza”. Piazzale A. Moro, 5, 00185 Roma, Italy.

²*Dipartimento di Matematica e Fisica, Università Roma Tre, Via della Vasca Navale 84, 00146 Roma, Italy.*

³*Istituto dei Sistemi Complessi - CNR, via del Fosso del Cavaliere 100, 00133 Rome, Italy.*

Langer theory of metastability provides a description of the lifetime and properties of the metastable phase of the Ising model field-driven transition, describing the magnetic field-driven transition in ferromagnets and the chemical potential-driven transition of fluids. An immediate further step is to apply it to the study of a transition driven by the temperature, as the one underwent by the two-dimensional Potts model. For this model a study based on the analytical continuation of the free energy (Meunier, Morel 2000) predicts the anomalous vanishing of the metastable temperature range in the limit of large system size, an issue that has been controversial since the eighties. With a parallel-GPU algorithm we compare the Monte Carlo dynamics with the theory, obtaining agreement and characterizing the anomalous system size dependence. We discuss the microscopic origin of these metastable phenomena, essentially different with respect to the Ising case.

Metastability is ubiquitous in nature and in technology [1]. It is an important concept in many fields of physics [2]. In particular, it is crucial in the context of the glass transition problem [1, 3, 4]. Metastability is also present in many biological systems, such as proteins [5] or nucleic acids [6]. Despite the relevance of the subject, there is no general theoretical framework allowing for the computation of the lifetime and properties of the Metastable Phase (MP) provided the details of the microscopic interaction [7]. On the other hand, understanding how finite-size effects can influence this peculiar state of matter is becoming more and more relevant with the increasing development of miniaturization processes and of nano- and bio-sciences [8–10].

Beyond mean-field approximation, metastable states correspond to *local* free energy saddle points in phase space and standard methods of statistical physics are not able to capture their properties [7]. The main theoretical tools to tackle them are based on restricted ensembles [11–14], which exclude phase-separated configurations from the partition function. Of particular relevance is the study by Langer [15, 16]. He showed that the MPs of the Ising/Lattice Gas Model (ILGM) can be described by the analytical continuation of the free energy f in the unstable phase, $\tilde{f}$. Its imaginary part, $\text{Im } \tilde{f}$, is in this context proportional to the nucleation rate, I , for a wide class of model dynamics [17]. Such an approach allows for the computation of $I(h, T)$ as a function of the under-critical temperature T and of the field $h > 0$, corresponding to the MP with negative magnetization [15–19]. The Langer description of the ILGM has been compared with dynamical methods [2], in which the MP is characterized as a stationarity of observables under a Markov-Chain Monte Carlo (MC) local dynamical update [20–23]. A general agreement between the Droplet Theory (DT) and dynamics is found in 2D, 3D. Finite-size effects are also well understood in the ILGM [21], and absent when the system linear size L is much larger

than the length scales involved in the nucleation process. While the relationship between $\text{Im } \tilde{f}$ and I has been proved for several systems, its general applicability is not known [2]. The analytical continuation of f *à la* Langer has been computed for the order-disorder, T -driven transition of the q -color Potts model (PM) in 2D [24]. The interest for this model lies in the fact that, at variance with the ILGM, for $q > 4$ it undergoes a discontinuous transition driven by the temperature, which represents the unique thermodynamic variable of the model. This essential difference with respect to the ILGM motivates the application of the Langer/Fisher methods. A further motivation to study the MPs is that their existence for large system area at inverse temperatures above the inverse transition temperature β_t still results unclear after a long debate. In this paper we shall investigate the effects of finite size on metastability in this paradigmatic model and settle, at least phenomenologically, the controversial about the disappearing of the MP in thermodynamic limit.

Binder first posed the question [25], relating it to that of the finiteness or divergence of the specific heat at β_t , c_t . This approach was recently pursued [26] with the help of Graphic Processing Units (GPUs), resulting in a finite c_t . The possible existence of a metastable interval results as well from pseudo-critical approaches, which find critical divergences at the (so called *pseudo-spinodal*) temperature $\beta_s > \beta_t$ [27–29], for different values of q . A different picture emerge from the study [24], the mentioned DT for the 2D $q > 4$ PM. The free energy f_a of the ensemble of droplets with area a is guessed, by compatibility with exact results at β_t , to be $\beta f_a = \delta \beta a - w_q a^{-2/3} - (7/3) \ln a$, being w_q the surface tension energy, inversely proportional to the correlation length ξ_q , and $\delta \beta = \beta - \beta_t$ the excess of inverse temperature. The resulting free energy of the disordered (stable) phase for $\delta \beta < 0$ is analytically continued for complex values of the inverse temperature, obtaining the function ϕ :

$$\phi(z) = \int_0^\infty \frac{d\zeta}{\zeta^3} e^{-\zeta} \left[e^{-z\zeta^{3/2}} - 1 - z\zeta^{3/2} \right] \quad (1)$$

where $z = -\delta\beta/w_q^{3/2} \in \mathbb{C}$. Then, the finite-size energy probability distribution (EPD) $P_{A,\delta\beta}(E)$ corresponding to a system with A sites at $\delta\beta = 0$ is obtained by a Laplace transform of ϕ , and extended to the MP at $\delta\beta > 0$ by re-weighting:

$$P_{A,0}(\varepsilon) = \frac{gAw_q^2}{2i\pi} \int_{-\infty}^{\infty} dz e^{w_q^2 A(\phi(z) - \varepsilon z)} \quad (2)$$

$$P_{A,\delta\beta}(\varepsilon) = P_{A,0}(\varepsilon) e^{-\delta\beta A\varepsilon},$$

g is a constant, ε is the energy per site and $\epsilon = (\varepsilon - \varepsilon^{(d)})/gw_q^{1/2}$ is a rescaled energy shifted by the average (disordered phase) energy, $\varepsilon^{(d)}$, at β_t [24]. $P_{A,0}$ happens to exhibit an anomalous slow dependence on the area, such that the metastable interval $[\beta_t : \beta_t + \beta^*(A)]$ in which the EPD presents a convex region shrinks to zero in the large- A limit. No metastability is found to exist for $A \rightarrow \infty$. The disappearance of the metastable interval for large A is also found in [31], by means of *MC dynamics* in the MP at small $\delta\beta > 0$. The metastable dynamics at $\delta\beta > 0$ has also been studied in [32], where an anomalous finite-size behavior has been pointed out.

There are several questions to be clarified. First, the apparent disagreement between the results of pseudo-critical attempts [27–29], indicating the existence of a metastable interval for large sizes, and those based on stability conditions of the MP [24], and on dynamics [31, 32]. Secondly, to what extent the results of Ref. [24] coincide with those of a dynamic sampling for $\delta\beta > 0$ [24, 31, 32]. Finally, the microscopic origin of the shrinking of the metastable interval for large sizes, which is absent in the ILGM paradigm. In this letter we perform a comparison (the first one, to our knowledge) between the results from the DT and the dynamical averages under a local MC updating rule. We report evidence of the *dynamical* metastable inverse temperature interval shrinking approximately as $\sim A^{-1/3}$. This scaling can be also derived from the analysis of [24] in saddle point approximation. A devoted GPU parallel algorithm has been developed for the efficient simulation of large sizes ($L \sim 1024$). We have also devised a method allowing for an accurate comparison with the DT [24], an unaccomplished task, to our knowledge. Let us describe our method. The q -PM [38] is defined in a configuration of A spins $\{\sigma_j\}_{j=1}^A$, $\sigma_j = 1, \dots, q$, with Hamiltonian $\mathcal{H} = -\frac{1}{2} \sum_{i,j} \delta_{\sigma_i, \sigma_j} \mathcal{A}_{ij}$, where $\mathcal{A}$ is the adjacency matrix, which corresponds to a 2D lattice with periodic boundary conditions. For $A \rightarrow \infty$, the model presents a first-order phase transition for $q > 4$ at an inverse temperature $\beta_t(q) = \ln(q^{1/2} + 1)$ [38]. Our analysis is

for $q = 12$, for which $\varepsilon^{(d)} \simeq -0.8637$ and $\xi_{12} \simeq 6.54$ lattice sites [39]. Starting from different random configurations, we perform series of Metropolis MC Markov chains which generate *sequences* of configurations $\{\sigma_t\}_t$ at different times, differing by $t_s = 128$ MC steps. For $\delta\beta > 0$, we compute averages from *sub-sequences of stationary configurations only*. As a criterion of stationarity, we impose that the temporal self-correlation function $C_t(t') = \sigma_t \cdot \sigma_{t+t'}$, that is the overlap between configurations at instants $t, t + t'$, does not differ too much with respect to $C_{t-\delta t}$, the same function $\delta t = 10^3$ MC steps before, the difference required to be of the order of its fluctuations in stable equilibrium [30]. We assume that the set of different sub-sequences of stationary configurations (with energy safely larger than the corresponding ordered equilibrium energy at $\delta\beta$) constitutes a “stationary ensemble” to be compared with the restricted ensemble approach implicit in the metastable EPD $P_{A,\delta\beta}$ of [24], since the convexity of the latter is expected to induce stationarity under a local dynamics. The adoption of such a stationary ensemble yields better results than the commonly used first-passage time method [31, 32], since it excludes non-equilibrium realizations with over-critical growing droplets [2, 7] and systematically discriminates fake energy plateaus [30]. Due to the low free-energy barriers of the model for moderate values of q , such a method is necessary for an accurate comparison with the theory [30]. Within such a stationary ensemble we measure the EPD. On the other hand, properties of the whole stationary sequence are averaged over different instances of the sequence. An example of such an average is the average time length of the stationary sub-sequence, that will be called the *lifetime of the MP*, $\tau(\delta\beta, A)$.

In Fig. 1 we present the EPD for $A = 256^2$, for different values of $\delta\beta$. Symbols are data from our simulation, while continuous lines are the prediction Eq. (2). We have calculated the last quantity by contour-integrating Eq. (2) through the steepest descent of the integral, the complex evaluation of ϕ requiring in turn the integration through the optimal contour in Eq. (1). The theoretical EPDs present a minimum at $\varepsilon_m(\delta\beta, A)$ (evidenced as the circle over the $\delta\beta = 0.0038$ curve), which increases with $\delta\beta$. Within Meunier and Morel theory, the MP is conceivable in the *restricted* range $\varepsilon > \varepsilon_m(\delta\beta, A)$, and the so called *pseudo-spinodal point*, $\beta^*(A)$, at which ε_m reaches the inflexion point $\varepsilon^*(A)$ (the vertical line in the figure) signals the endpoint of metastability (the figure indicates that $\delta\beta^*(256^2)$ is somewhere in between 0.0038 and 0.0048). For $\delta\beta < \delta\beta^*$, the agreement between theory and numerics is good. We attribute the progressive discrepancies for large $|\varepsilon|$ s to a statistical underestimation for low probabilities and to a finite-size effect, presumably consequence of the continuum description of Eq. (1). On the other hand, for $\delta\beta > \delta\beta^*$ the difference between theory and numerics becomes essential: while the theory predicts absence of metastability (no convex $P_{\delta\beta,A}$), we




FIG. 1. Logarithm of the EPD for different inverse temperatures $\delta\beta$ (on the key) for $A = 256^2$, vs. the theoretical prediction Eq. (2) (curves). Inset: the quantity $\ln P_{\delta\beta,A}(\epsilon) - \delta\beta A \epsilon$, for different $\delta\beta$ s from 0.0016 to 0.08. Only for low $\delta\beta \lesssim 0.0046$ the data overlap for different $\delta\beta$ s (they satisfy re-weighting).

nevertheless still observe stationary states with convex P . Moreover, as illustrated in the inset, the EPD curves for different values of $\delta\beta > \delta\beta^*$ are not derivable one from another by re-weighting (as for $\delta\beta < \delta\beta^*$ they do, see Eq. (3)). This is an evidence of the fact that these stationary states do not obey Maxwell-Boltzmann statistics. An interesting question is whether the lifetime of the non-equilibrium stationary states for $\delta\beta > \delta\beta^*$ remains finite in the large- A limit. This point will be discussed below.

At the pseudo-spinodal point $\delta\beta^*(A)$, the slope of the EPD at $\epsilon^*(A)$ vanishes. We have estimated it by extrapolating in $\delta\beta$ our data for $(\partial_{\epsilon}|_{\epsilon^*} \ln P)_{\delta\beta,A}$ down to zero, obtaining values of $\delta\beta^*(A)$ in reasonable (better the larger the size) agreement with those of the DT (see Fig. 3). The data approximately exhibit the size dependence $\delta\beta^*(A) = C + D A^{-1/3}$, with a small C . The $\delta\beta^* \sim A^{-1/3}$ scaling is compatible with the DT, as we find for $\delta\beta \searrow 0$. The numerical results for $\delta\beta < \delta\beta^*$ are, hence, compatible with the DT picture, which predicts absence of equilibrium metastability for large A .

A direct comparison with the theory is possible also for the lifetime $\tau(\delta\beta, A)$. We assume that the lifetime of the MP is the time needed to create a fluctuation with energy lower than the limit of stability $\epsilon_m(\delta\beta, A)$, i.e., $\tau/\tau_r \sim P_{\delta\beta,A}(\epsilon_m(\delta\beta, A))^{-1}$. τ is to be measured in units of the self-correlation time τ_r , since the probability of a critical configuration is inversely proportional to the average number of *uncorrelated* configurations before leaving the plateau. Using $P_{\delta\beta,A}(\epsilon_m)$ from the solution of Eq. (2) (and neglecting the $\delta\beta$ dependence of τ_r) we have estimated the β -dependence of τ , which agrees with our numerical results in the DT validity region $\delta\beta < \delta\beta^*(A)$.

The comparison is illustrated in Fig. 2, in which we also report the analytical expression for $\ln \tau$, that we have obtained in saddle-point approximation to Eq. (2), valid for $\delta\beta \searrow 0$ [30]:

$$\ln \tau(\delta\beta, A) \sim p \ln \frac{2w_q}{3\delta\beta} + \frac{4w_q^3}{27} \frac{1}{\delta\beta^2} + \mathcal{K}_A, \quad (3)$$

with $p = 7$, and $\mathcal{K}$ being a decreasing function of the area, constant in $\delta\beta$. Interestingly, assuming $\tau \sim I^{-1}$ with I given by the Langer relation $I \sim \text{Im} \phi(z \nearrow 0)$, leads alternatively for τ to the same law as in Eq. (3) with $p = 5$ (see Fig. 2), although this simpler approach does not provide information about $\mathcal{K}_A$. For fixed temperatures, the numerical data for τ systematically decrease with A , in agreement with the DT.

From our data of τ we estimate the *dynamical temperature* $\delta\beta_d(A, t)$ at which the average lifetime is t . Remarkably, also our results for $\delta\beta_d$ present a clear scaling:

$$\delta\beta_d(A, t) = C_t A^{-1/3} + D_t \quad (4)$$

with nonzero D_t (see Fig. 3). We define in this way a temperature endpoint of the stationary phase, defined by the temperature at which τ becomes small. Choosing $t_{\min} = 6 \cdot 10^3$ MC steps, we obtain a threshold, $\delta\beta_d(A, t_{\min})$, below which the stationary states are no longer observable in practice. Eq. (4) suggests that in the large- A limit there may be a nonzero temperature interval $[\beta_t : \beta_t + D_{t_{\min}}]$ in which the non-Boltzmann stationary states indeed survive with a nonzero lifetime.

Our findings are summarized in Fig. 3. For $\delta\beta < \delta\beta^*(A)$ the local dynamics leads to Boltzmann DT-describable metastable states. On the other hand, for $\delta\beta > \delta\beta^*$ and sufficiently low, we expect stationary states with non-Boltzmann statistics. Finally, for $\delta\beta \gtrsim \delta\beta_d(A, t_{\min})$, no stationary states, just off-equilibrium relaxation towards the ordered phase is observed.

The shrinking of the metastable interval may not be in contradiction with the results of the aforementioned pseudo-critical studies: the limit β_s , where τ_r is supposed to diverge, cannot be reached in a dynamical scheme, in which metastability is supposed to end at lower β when τ_r becomes of the order of τ . The argument $\tau(\delta\beta^*, A) \sim \tau_r(\delta\beta^*, A)$ is indeed compatible with the decreasing of $\delta\beta^*(A)$, since as A increases, τ decreases and τ_r anomalously increases [30]. This dynamical meaning of $\delta\beta^*$ is also compatible with our data, as we anticipate in Fig. 2.

An important point concerns the microscopic origin of these different behaviors, which remains unknown. In [32] it is conjectured that the canonical spinodal point is associated to the Evaporation-Condensation transition [33, 34], a finite-size effect occurring at coexistence in the micro-canonical ensemble, and that has been observed for

the PM in 2D (when the micro-canonical endpoint of the stable energy branch is again at a value $\delta\beta \sim A^{-1/3}$) [32, 35, 36] and 3D [37]. The authors of [32] also conjecture the existence of a length scale ℓ (suggested to be the typical distance between critical droplets, R_0) such that the condition $A \sim \ell^2$ would trigger a crossover between different dynamical regimes (see also [37]). In any case, in the so called *deterministic region*, $A \gg R_0^2$, finite-size effects should disappear, as happens in the ILGM [21]. The decreasing of τ with A for arbitrarily large A is essentially different with respect to the ILGM paradigm, and implies a size-dependent nucleation rate. In [31], it is proposed that, while the bulk term in f_a of the ILGM is size-independent, in the PM case it may come from an entropy-maximizing constraint, such that clusters of a given color are confined to avoid the breaking of the symmetry between colors. In larger systems, clusters would be less confined since they contribute less to the magnetization.

The DT cannot validate this idea since it is obtained from an infinite-volume free energy, not allowing for a microscopic formulation in terms of droplets. We are investigating an alternative scheme by estimating the form of the size-dependent droplet free-energy $f_{a,A}$. The dependence of the results on lattice, algorithm dynamics and q value are also being examined.

Summing up, we have provided a picture of the metastable dynamics of the 2D PM. The MPs present a strong finite-size scaling, well described by the DT. Such an anomalous finite-size scaling, whose microscopic origin is controversial, is different with respect to the metastability in the ILGM, and may be found in other first-order transitions.

Acknowledgments. We acknowledge Kurt Binder, Ezequiel E. Ferrero and Per Arne Rikvold for critical comments on the manuscript. We gratefully acknowledge the support of NVIDIA Corporation with the donation of the Tesla K20 GPU used for this research.




FIG. 2. Logarithm of the lifetime for $A = 128^2, 256^2$, vs. the DT prediction (full color curves) [30], along with Eq. (3) with $p = 5$ and 7 (see text), and $\mathcal{K}_A$ taken as the only adjustable parameter. The vertical strips signal the error intervals around $\delta\beta^*(A)$. For small $\delta\beta$ we show τ_r , the data suggest that $\tau_r \sim \tau$ at $\delta\beta^*$.




FIG. 3. Pseudo-spinodal point $\delta\beta^*(A)$ extrapolated from our data vs. the DT predictions (carrés); $\delta\beta_d(A, t)$ for two values of t and a fit with Eq. (4) (lines). We also report $\delta\beta_B(A)$ at which the EPD stops being reweightable. Three different dynamical finite-size regimes emerge (see main text).

- [1] P. Debenedetti, *Metastable Liquids: Concepts and Principles*, Physical chemistry : science and engineering (Princeton University Press, 1996) ISBN 9780691085951
- [2] P. A. Rikvold and B. M. Gorman, “Recent results on the decay of metastable phases,” in *Annual Reviews of Computational Physics I* (1995) Chap. 5, pp. 149–191, cond-mat/9407027 see also: C. C. A. Günther, P. A. Rikvold, and M. A. Novotny, Phys. Rev. Lett. **71**, 3898 (Dec 1993) C. Günther, P. Rikvold, and M. Novotny, Physica A: Statistical Mechanics and its Applications **212**, 194 (1994), ISSN 0378-4371
- [3] P. G. Debenedetti and F. H. Stillinger, Nature **410**, 259 (Mar. 2001), ISSN 0028-0836
- [4] A. Cavagna, Physics Reports **476**, 51 (2009), ISSN 0370-1573
- [5] D. Thirumalai and G. Reddy, Nature chemistry **3**, 910

- (2011)
- [6] J. Völker, H. H. Klump, and K. J. Breslauer, Proceedings of the National Academy of Sciences **105**, 18326 (2008)
- [7] K. Binder, Reports on Progress in Physics **50**, 783+ (Jul. 1987), ISSN 0034-4885
- [8] M. D. Uchic, D. M. Dimiduk, J. N. Florando, and W. D. Nix, Science **305**, 986 (2004)
- [9] L. Sandoval and H. M. Urbassek, Nano Letters **9**, 2290 (2009), pMID: 19438190
- [10] M. S. Islam, J. Kano, and S. Kojima, Ferroelectrics **402**, 16 (2010)
- [11] D. Capocaccia, M. Cassandro, and E. Olivieri, Communications in Mathematical Physics **39**, 185 (1974), ISSN 0010-3616

- [12] O. Penrose and J. Lebowitz, *Journal of Statistical Physics* **3**, 211 (1971), ISSN 0022-4715
- [13] F. H. Stillinger, *Phys. Rev. E* **52**, 4685 (Nov 1995)
- [14] D. S. Corti and P. G. Debenedetti, *Industrial & Engineering Chemistry Research* **34**, 3573 (1995)
- [15] J. Langer, *Annals of Physics* **41**, 108 (1967), ISSN 0003-4916
- [16] J. S. Langer, *Phys. Rev. Lett.* **21**, 973 (Sep 1968)
- [17] J. Langer, in *Systems Far from Equilibrium*, Lecture Notes in Physics, Vol. 132, edited by L. Garrido (Springer Berlin Heidelberg, 1980) pp. 12-47, ISBN 978-3-540-10251-9
- [18] N. J. Gunther, D. J. Wallace, and D. A. Nicole, *Journal of Physics A: Mathematical and General* **13**, 1755 (1980)
- [19] F. Schmitz, P. Virnau, and K. Binder, *Phys. Rev. E* **87**, 053302 (May 2013)
- [20] K. Binder and H. Müller-Krumbhaar, *Phys. Rev. B* **9**, 2328 (Mar 1974)
- [21] P. A. Rikvold, H. Tomita, S. Miyashita, and S. W. Sides, *Phys. Rev. E* **49**, 5080 (Jun 1994)
- [22] K. Binder and E. Stoll, *Phys. Rev. Lett.* **31**, 47 (Jul 1973)
- [23] D. Heermann, A. Coniglio, W. Klein, and D. Stauffer, *Journal of Statistical Physics* **36**, 447 (1984), ISSN 0022-4715
- [24] J. L. Meunier and A. Morel, *The European Physical Journal B - Condensed Matter and Complex Systems* **13**, 341 (Jan. 2000), ISSN 1434-6028
- [25] K. Binder, *Journal of Statistical Physics* **24**, 69 (Jan. 1981), ISSN 0022-4715
- [26] E. E. Ferrero, J. P. D. Francesco, N. Wolovick, and S. A. Cannas, *Computer Physics Communications* **183**, 1578 (2012), ISSN 0010-4655
- [27] L. Fernández, J. Ruíz-Lorenzo, M. Lombardo, and A. Tarancón, *Physics Letters B* **277**, 485 (1992), ISSN 0370-2693
- [28] L. Schülke and B. Zheng, *Phys. Rev. E* **62**, 7482 (Nov 2000)
- [29] E. S. Loscar, E. E. Ferrero, T. S. Grigera, and S. A. Cannas, *The Journal of Chemical Physics* **131**, 024120 (2009)
- [30] M. I. Berganza, P. Coletti, and A. Petri, in preparation(2014)
- [31] M. Ibáñez Berganza, A. Petri, and P. Coletti, *ArXiv e-prints*(Jun. 2007), arXiv:0706.3534 [cond-mat.stat-mech]
- [32] T. Nogawa, N. Ito, and H. Watanabe, *Physics Procedia* **15**, 76 (2011), ISSN 1875-3892, proceedings of the 24th Workshop on Computer Simulation Studies in Condensed Matter Physics (CSP2011)
- [33] K. Binder and M. Kalos, *Journal of Statistical Physics* **22**, 363 (1980), ISSN 0022-4715
- [34] K. Binder, *Physica A: Statistical Mechanics and its Applications* **319**, 99 (2003), ISSN 0378-4371
- [35] T. Nogawa, N. Ito, and H. Watanabe, *Phys. Rev. E* **84**, 061107 (Dec 2011)
- [36] W. Janke, *Nuclear Physics B - Proceedings Supplements* **63**, 631 (Apr. 1998), ISSN 09205632
- [37] A. Bazavov, B. A. Berg, and S. Dubey, *Nuclear Physics B* **802**, 421 (2008), ISSN 0550-3213
- [38] F. Y. Wu, *Reviews of Modern Physics* **54**, 235 (Jan. 1982)
- [39] C. Borgs and W. Janke, *Journal de Physique I* **2**, 2011 (Nov. 1992)